

09/828,211

Generate Collection

Print

**Search Results - Record(s) 1 through 3 of 3 returned.**

- ☐ 1. 6287822. 04 Aug 98; 11 Sep 01. Mutation detection method. Gjerde; Douglas T, et al. 435/91.2; 210/198.2 210/635 435/6 536/22.1 536/25.4. C12P019/34 C12Q001/68 C07H019/00 C07H021/04 B01D015/08.
- ☐ 2. 5795976. 08 Aug 95; 18 Aug 98. Detection of nucleic acid heteroduplex molecules by denaturing high-performance liquid chromatography and methods for comparative sequencing. Oefner; Peter Josef, et al. 536/25.4; 204/450 204/456 435/6 435/91.2 536/22.1 536/23.1 536/24.3. C12N015/10 C07H021/02 C12Q001/68 C12P019/34.
- ☐ 3. 4879214. 15 Nov 88; 07 Nov 89. Differentiation of nucleic acid segments on the basis of nucleotide differences. Kornher; John S., et al. 435/6; 435/91.2. C12Q001/68.

Generate Collection

Print

Term	Documents
LABEL\$2	0
LABEL.DWPI,EPAB,JPAB,USPT.	110811
LABELA.DWPI,EPAB,JPAB,USPT.	3
LABELAD.DWPI,EPAB,JPAB,USPT.	1
LABELB.DWPI,EPAB,JPAB,USPT.	1
LABELBY.DWPI,EPAB,JPAB,USPT.	1
LABELC.DWPI,EPAB,JPAB,USPT.	1
LABELCC.DWPI,EPAB,JPAB,USPT.	1
LABELCD.DWPI,EPAB,JPAB,USPT.	1
LABELCO.DWPI,EPAB,JPAB,USPT.	1
LABELD.DWPI,EPAB,JPAB,USPT.	30
(L3 AND LABEL\$2).USPT,JPAB,EPAB,DWPI.	3

There are more results than shown above. [Click here to view the entire set.](#)

[Previous Page](#)[Next Page](#)

**WEST****Freeform Search**

Database:

US Patents Full-Text Database	▲
US Pre-Grant Publication Full-Text Database	
JPO Abstracts Database	
EPO Abstracts Database	
Derwent World Patents Index	
IBM Technical Disclosure Bulletins	▼

Term:

L3 and label\$2	▲
	▼

Display:  Documents in Display Format:  Starting with Number

Generate: ☐ Hit List ☒ Hit Count ☐ Side by Side ☐ Image

Search	Clear	Help	Logout	Interrupt
--------	-------	------	--------	-----------

Main Menu	Show S Numbers	Edit S Numbers	Preferences	Cases
-----------	----------------	----------------	-------------	-------

**Search History**

DATE: Tuesday, April 16, 2002   [Printable Copy](#)   [Create Case](#)

Set Name   Query  
side by side

Hit Count   Set Name  
result set

DB=USPT,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ

<u>L4</u>	L3 and label\$2	3	<u>L4</u>
<u>L3</u>	L2 and heteroduplex\$2 and homoduplex\$2	6	<u>L3</u>
<u>L2</u>	L1 and (reverse phase near5 chromatograph\$)	39	<u>L2</u>
<u>L1</u>	(detect\$ or determin\$) near5 mismatch	3569	<u>L1</u>

END OF SEARCH HISTORY

## End of Result Set



Generate Collection

L4: Entry 3 of 3

File: USPT

Nov 7, 1989

DOCUMENT-IDENTIFIER: US 4879214 A

TITLE: Differentiation of nucleic acid segments on the basis of nucleotide differences

Brief Summary Paragraph Right (7):

A technique for detecting specific mutations in any segment of DNA is described in Wallace, et al., Nucl. Acids Res. 9:879-894 (1981). It involves hybridizing the DNA to be analyzed (target DNA) with a complementary, labeled oligonucleotide probe. Due to the thermal instability of DNA duplexes containing even a single base pair mismatch, differential melting temperature can be used to distinguish target DNAs that are perfectly complementary to the probe from target DNAs that differ by as little as a single nucleotide. An adaptation of this technique, described by Saiki, et al., U.S. Pat. No. 4,683,194, can be used to detect the presence or absence of a specific restriction site. In Saiki's adaptation, an end-labeled oligonucleotide probe spanning a restriction site is hybridized to the target DNA. The hybridized duplex of DNA is then appropriately incubated with the restriction enzyme for that site. Only paired duplexes between probe and target that reform the restriction site will be cleaved by digestion with the restriction endonuclease. Detection of shortened probe molecules indicates that the specific restriction site is present in the target DNA. In a related technique, described in Landegren, et al., Science 241:1077-1080 (1988), oligonucleotide probes are constructed in pairs such that their junction corresponds to the site on the DNA being analyzed for mutation. These oligonucleotides are then hybridized to the DNA being analyzed. Base pair mismatch between either oligonucleotide and the target DNA at the junction location prevents the efficient joining of the two oligonucleotide probes by DNA ligase. A major problem with these and other oligonucleotide techniques is that the mutation must already be characterized as to type and location in order to synthesize the proper probe. Thus, techniques using oligonucleotide probes can be used to assay for specific, known mutations, but they cannot be used generally to identify previously undetected mutations.

Brief Summary Paragraph Right (8):

In the technique described in Mundy, U.S. Pat. No. 4,656,127, specific mutations can be detected by first hybridizing a labeled DNA probe to the target nucleic acid in order to form a hybrid in which the 3' end of the probe is positioned adjacent to the specific base being analyzed. Then, a DNA polymerase is used to add a nucleotide analog, such as a thionucleotide, to the probe strand, but only if the analog is complementary to the specific base being analyzed. Finally, the probe-target hybrid is treated with exonuclease III. If the nucleotide analog has been incorporated, the labeled probe is protected from nuclease digestion. Absence of a labeled probe indicates that the analog and the specific base being analyzed were not complementary. As with abovediscussed techniques involving oligonucleotides, this method detects specific mutations, but it cannot be used in a general manner to detect all possible nucleotide differences.

Brief Summary Paragraph Right (9):

Nucleotide differences between two DNA sequences also can be studied by forming a heteroduplex between the two DNAs of interest. Base pair mismatches will occur within the heteroduplex at points where the sequences differ. A number of methods have been developed to detect such mismatches. Chemical probes for mismatches exist that specifically react with those atoms in the base normally involved in hydrogen bonding, see, e.g., Novack, et al., Proc. Natl. Acad. Sci. U.S.A. 83:586-590 (1986); Cotton, et al., Proc. Natl. Acad. Sci. U.S.A. 85:4397-4401 (1988). These chemically altered sites are susceptible to chemical cleavage, whereas a perfectly paired duplex is not. Problems with this technique include: (i) toxicity of the chemical reagents and (ii)

efficiency of much less than 100% for the reactions with unpaired bases. Another approach to mismatch detection is based upon the ability of certain nucleases to recognize and cleave these sites. S.sub.1 nuclease and RNase A have been shown effective in mismatch detection and cleavage, see, e.g., Shenk, et al., Proc. Natl. Acad. Sci. U.S.A. 72:989-993 (1975); Myers, et al., Science 230:1242-1246 (1985). Neither of these enzymes, however, cleaves at all possible mismatched base pairs. There is also considerable background associated with nuclease cleavage at perfectly paired sites in the duplex.

Brief Summary Paragraph Right (10):

Myers, et al., Nature 313:495-498 (1985), and Fischer, et al., Proc. Natl. Acad. Sci. U.S.A. 80:1579-1583 (1983), have demonstrated that mismatched base pairs within a heteroduplex alter its melting properties with respect to a perfectly paired homoduplex. These altered melting properties can be observed electrophoretically on a gel containing an exponential gradient of denaturant. A serious drawback to this technique is the difficulty of manipulating and processing these gels. Another problem is the inability of this technique to detect uniformly all base pair mismatches along a given heteroduplex. The resolving power of these gels is reduced with increasing GC content of the heteroduplex, and thus mutations in a GC-rich domain are more difficult to detect than mutations in a domain with a lower GC content.

Brief Summary Paragraph Right (11):

The primer extension process described in Proudfoot, et al., Science 209:1329-1336 (1980), has been widely used to study the structure of RNA and also has been used to characterize DNA, see, e.g. Engelke, et al., Proc. Natl. Acad. Sci. U.S.A. 85:544-548 (1988). This process consists of hybridizing a labeled oligonucleotide primer to a template RNA or DNA and then using a DNA polymerase and deoxynucleoside triphosphates to extend the primer to the 5' end of the template. The labeled primer extension product is then fractionated on the basis of size, usually by electrophoresis through a denaturing polyacrylamide gel. When used to compare homologous DNA segments, this process can detect differences due to nucleotide insertion or deletion. Because size is the sole criterion used to characterize the primer extension product, this method cannot detect differences due to nucleotide substitution.

Brief Summary Paragraph Right (39):

In order to compare the mobility of the synthesized strands, the strands must be detected. Any method used to detect nucleic acid can be used to detect the synthesized strands containing mobility-shifting nucleotide analogs. One method is to perform the primer extension reaction with a labeled primer. Typically, oligonucleotide primers are labeled by attaching a reporter to the 5' end of the oligonucleotide. This reporter can be a radioactive isotope; a group that takes part in an enzyme or fluorescent or chemiluminescent reaction; or some other small molecule, such as biotin, that can be detected by a convenient method. Alternatively, labeling can be performed by incorporating reporter-tagged nucleotides into the synthesized strands as they are synthesized. The reporter can be attached either to a mobility-shifting nucleotide analog or to one of the other nucleotides used as substrates in the preparation of synthesized strands. The same type of reporters used to label primers can be used to label the nucleotide substrates. Another method of detection entails hybridizing a reporter-labeled probe to the synthesized strands after the size fractionation step, as in the Southern blot procedure.

Detailed Description Paragraph Right (6):

Next, the triethylammonium salt of 5-(3-biotinamido(hexanoylamido))-1-propynyl)-2'-deoxycytidine 5'-triphosphate, 5-(Bio-AC-AP3)dCTP, ##STR5## was prepared from 5-(3-amino-1-propynyl)-2'-deoxycytidine 5'-triphosphate, 5-(AP3)dCTP. 5-(3-Amino-1-propynyl)-2'-deoxycytidine 5'-triphosphate (30  $\mu$ mol) was dissolved in 1 M aqueous TEAB, pH 7.6, (600  $\mu$ L) and sulfosuccinimidyl 6-(biotinamido)-hexanoate Na salt (Pierce Chemical Co., Rockford, IL; 33 mg; 60  $\mu$ mol) was added. The solution was held at 50.degree. C. for 90 min and then diluted to 6 mL with water. The solution was loaded onto a DEAE Sephadex A-25-120 column (1.times.19 cm) that had been equilibrated with 1.0 M aqueous TEAB, pH 7.6. The column was eluted with a linear gradient of TEAB, pH 7.6, from 0.1 M (150 mL) to 1.0 M (150 mL), running at c. 100 mL/hr, collecting fractions every 3 min. The eluent was monitored by absorbance at 270 nm, and the fractions corresponding to the second peak (#22-33) were pooled, stripped, and co-evaporated (2x) with ethanol. The residue was

subjected to reverse phase chromatography (Baker 7025-00) on a column (1.times.8 cm) poured in acetonitrile/1 M aqueous TEAB, pH 7.6. The column was eluted with a step gradient of acetonitrile/1 M TEAB, pH 7.6, (0-30% acetonitrile, 2%/step, 2 mL/step, 1 fraction/step). The fractions were assayed by HPLC on a reverse phase column under conditions analogous to those used in the above separation. Starting material eluted first (#2-6) followed by the product (#7-11). The product fractions were pooled, stripped, and coevaporated (2x) with ethanol. The yield, assuming an absorption coefficient of 10,100 at 295.5 nm, was 10.5  $\mu$ mol (35%). HPLC on Zorbax SAX and on reverse phase showed the material to be >99% pure. The .sup.1 H- and .sup.31 P-NMR spectra are fully consistent with the title structure.

☐ Generate Collection

L4: Entry 2 of 3

File: USPT

Aug 18, 1998

DOCUMENT-IDENTIFIER: US 5795976 A

TITLE: Detection of nucleic acid heteroduplex molecules by denaturing high-performance liquid chromatography and methods for comparative sequencingAbstract Paragraph Left (1):

The present invention describes a method for separating heteroduplex and homoduplex DNA molecules in a mixture. In the method, such a mixture is applied to a stationary reverse phase support. The heteroduplex and homoduplex molecules are eluted with a mobile phase containing an ion-pairing reagent and an organic solvent. The eluting is carried out under conditions effective to at least partially denature the heteroduplexes (e.g., thermal or chemical denaturing) resulting in the separation of the heteroduplexes from the homoduplexes. The method has many applications including, but not limited to, comparative nucleic acid sequencing, linkage analysis, evolutionary studies, forensics, identification of disease-causing gene mutations, genetic marker development and diagnostics.

Brief Summary Paragraph Right (1):

The present invention relates to a chromatographic method for detecting heteroduplexes in nucleic acid fragments, and particularly to denaturing high performance liquid chromatography, for use in detecting mutations and for comparative DNA sequencing.

Brief Summary Paragraph Right (11):

Presently used methods for screening for polymorphic sites within a gene include RNase A cleavage, chemical cleavage, denaturing gradient gel electrophoresis. These methods exploit characteristics of mismatched heteroduplexes formed between normal and mutant sequences.

Brief Summary Paragraph Right (14):

In denaturing gradient gel electrophoresis (DGGE), either homoduplex or heteroduplex double stranded DNA is electrophoresed under denaturing conditions of increasing concentration until the last domain is denatured, and migration of the DNA through the gel is retarded. DNA sequences differing by a single base pair migrate at different rates along the gel, thereby allowing detection of a polymorphic site, if present.

Brief Summary Paragraph Right (17):

The ligase-mediated method for detecting mutations makes use of the fact that the ends of two single strands of DNA must be exactly aligned for DNA ligase to join them. In utilizing this technique, oligonucleotides complementary to the target sequence, 5' to and including the mutation site, are synthesized and labeled. A third oligonucleotide complementary to the common sequence 3' to the mutation site is synthesized and also labeled. The oligonucleotides are then hybridized to strands of the target. In cases in which the 5' and 3' oligonucleotides form a flush junction that can be joined by DNA ligase, ligation occurs. However, a single base pair mismatch occurring between the normal 5' oligonucleotide and the mutation site is sufficient to prevent the ligase from acting and can readily be detected.

Brief Summary Paragraph Right (19):

The present invention provides a method for separating heteroduplex and homoduplex nucleic acid molecules (e.g., DNA or RNA) in a mixture using high performance liquid chromatography. In the separation method, a mixture containing both heteroduplex and homoduplex nucleic acid molecules is applied to a stationary reverse-phase support. The sample mixture is then eluted with a mobile phase containing an ion-pairing reagent and an organic solvent. Sample elution is carried out under conditions effective to at least partially denature the heteroduplexes and results in the separation of the heteroduplex and homoduplex molecules.

Brief Summary Paragraph Right (23):

At least partial denaturation of heteroduplex molecules can be carried out several ways including the following. Temperatures for carrying out the separation method of the invention are typically between about 50.degree. and 70.degree. C., preferably between about 55.degree.-65.degree. C. In a preferred embodiment, the separation is carried out at 56.degree. C. Alternatively, in carrying out a separation of GC-rich heteroduplex and homoduplex molecules, a higher temperature (e.g., 64.degree. C.) is preferred.

Brief Summary Paragraph Right (24):

Alternately, sample elution is carried out under pH conditions effective to at least partially denature the heteroduplex molecules. In such cases, a lower column temperature less than about 65.degree. C. may be sufficient for the separation of the heteroduplex and homoduplexes molecules in the sample.

Brief Summary Paragraph Right (26):

In one particular embodiment of the present method, homoduplex and heteroduplex molecules in a mixture are separated by applying the mixture to a C-18 alkylated polystyrene-divinylbenzene copolymer stationary support and eluting the mixture with a mobile phase containing triethylammonium acetate as the ion-pairing reagent and acetonitrile as the organic solvent at a temperature between about 50.degree.-65.degree. C.

Brief Summary Paragraph Right (27):

In an alternate embodiment, the homoduplex and heteroduplex molecules contained in the mixture are amplified using the polymerase chain reaction and the amplified DNA molecules are denatured and renatured to form a mixture of heteroduplex and homoduplex molecules prior to carrying out the separation method of the invention.

Brief Summary Paragraph Right (30):

Also disclosed is a method for comparative DNA sequencing in which potentially all possible nucleotide mismatches and insertion/deletions within select amplified DNA fragments obtained from multiple animal or human subjects can be detected. In the context of the present invention, comparative DNA sequencing is carried out by amplifying DNA samples, typically up to at least about 1.5 kb in length, obtained from multiple subjects. The amplified DNA fragments are then surveyed, either individually or in pools containing up to about 10 samples, for the presence or absence of heteroduplexes using the denaturing high performance liquid chromatography method of the present invention.

Brief Summary Paragraph Right (31):

In surveying the samples, the amplified DNA fragments are denatured and allowed to reanneal. The resulting mixture of DNA fragments is then applied to a stationary reverse-phase support. The sample mixture is eluted with a mobile phase containing an ion-pairing reagent and an organic solvent. Sample elution is carried out under conditions effective to at least partially denature any heteroduplexes present in the sample and results in the detection of any heteroduplex molecules contained in the sample. The detection of a heteroduplex indicates the presence of a base pair mismatch and/or an insertion/deletion in the sample fragment(s).

Brief Summary Paragraph Right (32):

In instances in which only homoduplexes are observed during the sample screening, further standard sequencing is not required since the sequence is monomorphic (i.e., lacking a polymorphic site) in all subjects compared. In utilizing the method of the present invention, only those DNA fragments identified as heteroduplexes, and therefore identified as containing at least one polymorphic site, are then sequenced by conventional methods to characterize the observed polymorphism(s).

Drawing Description Paragraph Right (5):

FIG. 5 is a scheme illustrating the mixing of two different double-stranded DNA fragments, subsequent denaturing, and reannealing to produce both homoduplex (2 species) and heteroduplex (2 species) products;

Drawing Description Paragraph Right (6):

FIGS. 6A-6C are IP-RP-HPLC chromatograms illustrating the detection at 50.degree. C.

of a 209-base pair homoduplex "homo-A-209" (FIG. 6A), a 209-base pair homoduplex "homo-G-209" (FIG. 6B), and a chromatogram of the hybrids formed after mixing homoduplexes A and G, denaturing, and reannealing to produce both homoduplex and heteroduplex products (FIG. 6C);

Drawing Description Paragraph Right (8):

FIGS. 8A-8C are IP-RP-HPLC chromatograms illustrating the detection (at 56.degree. C.) of two 439 base pair heteroduplexes each containing a single site mismatch in a sample containing both 439-mer homoduplex and heteroduplex samples (FIG. 8C). FIGS. 8A and 8B are chromatograms of each of the two different 439-mer homoduplexes designated as "homo-A-439" and "homo-G-439", respectively, prior to mixing, denaturing, and reannealing;

Drawing Description Paragraph Right (9):

FIG. 9A-9C are IP-RP-HPLC chromatograms of each of two PCR-amplified 1-kilobase homoduplexes, "homo-A-1kb" (FIG. 9A) and "homo-G-1kb" (FIG. 9B). FIG. 9C illustrates the detection, at 56.degree. C., of two 1-kilobase heteroduplexes, each containing a single base-pair mismatch, in a mixture also containing two 1-kilobase homoduplexes;

Drawing Description Paragraph Right (10):

FIG. 10-10C are IP-RP-HPLC chromatograms illustrating the separation of 209-base pair homoduplexes from 209-base pair heteroduplexes as a function of column temperature;

Drawing Description Paragraph Right (12):

FIGS. 12A-C are IP-RP-HPLC chromatograms of each of two PCR-amplified 1.5-kilobase homoduplexes (FIGS. 12A and 12B) and the detection, at pH 7.5, of two 1.5-kilobase heteroduplexes, each containing a single base-pair mismatch (FIG. 12C); and

Drawing Description Paragraph Right (13):

FIGS. 13A-13E are IP-RP-HPLC chromatograms of a segment of the .beta.-globin gene containing a polymorphic site flanked by a GC-rich region, illustrating the effect of temperature on sample resolution for detecting heteroduplexes.

Detailed Description Paragraph Right (3):

"Ion-pair (IP) chromatography" as used herein refers to any chromatographic method for separating samples in which some or all of the sample components contain functional groups which are ionized or are ionizable. Ion-pair chromatography is typically carried out with a reverse phase column in the presence of an ion-pairing reagent.

Detailed Description Paragraph Right (7):

"Ion pairing reverse phase high performance liquid chromatography (IP-RP-HPLC)" refers to a type of high performance liquid chromatography in which the solid support is a reverse phase support and the mobile phase contains an ion-pairing reagent.

Detailed Description Paragraph Right (10):

"Homoduplex molecules" are typically composed of two complementary DNA strands.

Detailed Description Paragraph Right (11):

"Heteroduplex molecules" are typically composed of two complementary nucleic acid strands (e.g., DNA or RNA), where the strands have less than 100% sequence complementarity. The functional definition of homoduplex and heteroduplex molecules, in the context of the present invention, is apparent from the results presented below. Typically, in a mixed population of homoduplex and heteroduplex molecules, for shorter strands (e.g., typically about less than 70 base pairs in size) heteroduplex molecules elute as peaks corresponding to their respective denatured single strands under select denaturing conditions using reverse phase ion pairing HPLC, separable from those of homoduplex molecules. In a mixed population of homoduplex and heteroduplex molecules larger than about 70 base pairs in length, heteroduplex molecules typically elute with shorter retention times than those of homoduplexes of essentially the same size under select denaturing conditions using reverse phase ion pairing HPLC. "Insertions or deletions" or "indels" can occur in duplexes consisting of two complementary DNA strands, where the first strand of the DNA contains a greater number of nucleotides at an internal site than the second strand DNA molecule, and where these extra nucleotides are flanked by paired-complementary sequences. Indels can occur in heteroduplexes.



Detailed Description Paragraph Right (12):

"Base-pair mismatches" typically refers to a single base-pair mismatch flanked by matched base-pairs. Base-pair mismatches also include a series of mismatched base-pairs flanked by matched base-pairs. Base-pair mismatches can occur in heteroduplexes.

Detailed Description Paragraph Right (13):

A heteroduplex molecule that is "at least partially denatured" under a given set of chromatographic conditions refers to a molecule in which several complementary base pairs of the duplex are not hydrogen-bond paired, such denaturing typically extending beyond the site of the base-pair mismatch contained in the heteroduplex, thereby enabling the heteroduplex to be distinguishable from a homoduplex molecule of essentially the same size. In accordance with the present invention, such denaturing conditions may be either chemically (e.g., resulting from pH conditions) or temperature-induced, or may be the result of both chemical and temperature factors.

Detailed Description Paragraph Right (14):

"Comparative DNA sequencing" as used herein refers to a method for detecting any, and preferentially all, possible nucleotide mismatches and insertion/deletions within select amplified or non-amplified DNA fragments obtained from multiple animal or human subjects. In comparative DNA sequencing, DNA samples, typically up to at least about 1.5 kb in length, are obtained from multiple subjects and amplified or are otherwise produced (e.g., by cloning). The amplified DNA fragments are then surveyed, either individually or in pools containing up to about 10 unique samples, for the presence or absence of heteroduplexes. The detection of a heteroduplex indicates the presence of a base pair mismatch and/or an insertion/deletion in the sample fragment(s). Although comparative sequencing can be carried out using any of a number of analytical methods, it is particularly suited to the denaturing high performance liquid chromatography method of the present invention.

Detailed Description Paragraph Right (15):

In the context of the present invention, in instances in which only homoduplexes are observed during the sample screening, further standard sequencing is not required since the sequence is monomorphic in all subjects compared. In utilizing the method of the present invention, only those DNA fragments identified as heteroduplexes, and therefore identified as containing at least one polymorphic site, are sequenced by conventional methods to characterize the observed polymorphism(s).

Detailed Description Paragraph Right (18):

The present invention provides a method for separating heteroduplex and homoduplex DNA molecules in a mixture using high performance liquid chromatography and more particularly, denaturing high performance liquid chromatography, as will be described in detail below. The method can be utilized for detecting a single base mismatch in a DNA duplex containing up to about 2000 base pairs.

Detailed Description Paragraph Right (19):

High performance liquid chromatography (HPLC) generally refers to a technique for partitioning a sample or more specifically the components of a sample between a liquid moving or mobile phase and a solid stationary phase. In the present invention, the applicants have discovered a chromatographic method which utilizes conditions effective for at least partially denaturing heteroduplexes during sample elution to thereby enable the separation and identification of heteroduplexes and homoduplexes contained in a mixture.

Detailed Description Paragraph Right (20):

In the method of the present invention, a sample mixture containing both heteroduplex and homoduplex molecules is applied to a stationary phase. Generally, the stationary phase is a reverse phase material in which the chemically bonded phase is hydrophobic and is less polar than the starting mobile phase. Any of a number of commercially available reverse phase solid supports may be utilized in the present nucleic acid separation method although the resolution may vary depending upon the nature of the sample and other relevant experimental parameters.

Detailed Description Paragraph Right (28):

For achieving separations of samples containing heteroduplexes and homoduplexes of up to about 2000 base pairs in size, the stationary phase will typically have a surface area of about 2-400 m.<sup>2</sup>/g, and preferably about 8-20 m.<sup>2</sup>/g as determined by nitrogen adsorption.

Detailed Description Paragraph Right (36):

In an alternate embodiment, the pH of the mobile phase is adjusted to effect at least partial denaturation of the heteroduplex molecules in a sample containing a mixture of homoduplexes and heteroduplexes to allow separation and detection of the heteroduplex molecules. In using chemical means to effect heteroduplex denaturation, the pH may be adjusted by addition of either base (e.g., sodium hydroxide or urea to a pH of around about 8) or acid (e.g., triethylamine and acetic acid at a pH of about 8) under conditions effective to at least partially denature the heteroduplex molecules and which do not degrade the nucleic acids present in the sample nor adversely affect the integrity of the stationary phase. In such cases, sample elution may be carried out at temperatures less than about 50.degree. C.

Detailed Description Paragraph Right (39):

One way to achieve the denaturing HPLC conditions of the present invention (e.g., effective to at least partially denature heteroduplexes) is to modulate column temperature, as will be discussed in reference to the Examples below.

Detailed Description Paragraph Right (40):

A column temperature typically between about 50.degree.-65.degree. C. is preferred for resolving heteroduplex molecules from their corresponding homoduplex molecules by denaturing HPLC chromatography. The optimal column temperature will depend upon the sequence (base composition) of the sample to be separated, the choice of stationary phase, the choice of mobile phase, pH, flow rate, and the like, and in many cases, will be determined empirically. Ideally, in cases with known sequence, a suitable column temperature may be calculated.

Detailed Description Paragraph Right (41):

As will be seen from the discussion of the Examples below, heteroduplex detection can, in many cases, be accomplished at a column temperature of about 56.degree. C. Column temperatures as low as about 50.degree. C. can be used for detecting point mutations in small nucleic acid fragments containing up to about 70 base pairs. Column temperatures up to about 63.degree. C. may be optimal for separating G-C rich fragments.

Detailed Description Paragraph Right (42):

In addition to providing a denaturing HPLC method for detecting heteroduplex molecules in a sample containing both heteroduplex and homoduplex molecules of essentially the same size, the present method can also be used to detect homoduplex molecules containing short single-stranded overhangs, as illustrated in Example 2. Further, by varying the column temperature, both partial and complete denaturation of a 30-mer homoduplex containing a 2-base overhang was detectable in the resulting chromatograms.

Detailed Description Paragraph Right (45):

In contrast, elution at 50.degree. C. was effective to denature all of the D-E hybrid molecules in sample, as seen in FIG. 2C. The two peaks in the resulting chromatogram correspond to single stranded oligonucleotides D and E, respectively. The above exemplary results demonstrate that under the denaturing HPLC conditions of the present invention, complementary hybrid molecules containing short overhangs can be denatured and that the extent (e.g., percentage of molecules denatured) of denaturation is a function of temperature. Utilizing the denaturing conditions of the present invention, short homoduplex fragments (i.e., less than about 70 base pairs in length) carrying short overhangs may be detected due to their denaturation during sample elution and detected in the resulting chromatograms as their corresponding single-stranded components.

Detailed Description Paragraph Right (46):

In initial experiments to test the general applicability of the method, a short heteroduplex 43-mer containing a single base mismatch and lacking an overhang was eluted using the denaturing conditions of the present invention to investigate whether

or not such conditions would be effective in denaturing the heteroduplex. The details of the experiment are described in Example 3.

Detailed Description Paragraph Right (47):

Briefly, liquid hybridization of two 43-mers, oligonucleotide C and oligonucleotide A, a 43-mer complementary, with the exception of one base located ten bases from the 5' end of oligo A, to oligonucleotide C, was carried out by heating an equimolar mixture of the two subject oligonucleotides. The resulting A-C heteroduplex containing a single base pair mismatch was then chromatographed on a C18-polystyrene-divinylbenzene support using a binary gradient system composed of 0.1 TEAA (triethylammonium acetate) and 0.1 TEAA containing 25% acetonitrile at two different temperatures, 40.degree. C. and 51.degree. C., respectively. Details of the method including the gradient profile used to elute the sample are provided in Example 3.

Detailed Description Paragraph Right (48):

The results of the above described chromatography are shown in FIGS. 3A-C and 4A-C. In contrast to the results described above for the 30-mer homoduplex with a 2-base overhang, at the lower operating temperature of 40.degree. C. (FIGS. 3A-C), elution of the A-C hybrid resulted in a chromatogram corresponding to the annealed oligonucleotide fragments (heteroduplex A-C), as indicated in the resulting chromatogram by a single peak with a retention time differing from that of either oligo A or C. Upon raising the column temperature to 51.degree. C., complete denaturation of the sample was observed, as indicated in FIGS. 4A-C by the detection of two single peaks with retention times corresponding to those of single stranded oligonucleotide A and single stranded oligonucleotide C, respectively. In the exemplary chromatography described above, the effect of increasing column temperature on denaturation and subsequent detection of a heteroduplex molecule under a given set of chromatographic conditions is shown. The above results also demonstrate the ability of the method to discriminate small heteroduplex molecules in a sample by denaturing HPLC and indicate a beneficial feature of the method, namely, to separate oligonucleotides not only as a function of their size but also as a function of their respective sequence.

Detailed Description Paragraph Right (49):

In further support of the method of the present invention, experiments have been carried out in which single base mismatches in heteroduplexes to 1.5 kilobase in size have been detected, as will be described below. Further, a single set of chromatographic conditions was effective for detecting single base mismatches in mixtures of homoduplex and heteroduplex molecules from about 200-1500 base pairs in size, demonstrating the general applicability of the method.

Detailed Description Paragraph Right (50):

In an exemplary method for detecting polymorphisms in nucleic acid fragments, experiments were carried out in which two different homoduplex molecules, identical in sequence with the exception of one base pair, were denatured and reannealed to form a mixture containing four resulting duplex products, two homoduplexes and two heteroduplexes. Experiments were carried out with pairs of homoduplexes as described above containing 209, 439, 1000, and 1500 base pairs, respectively. Details of the experiments are described in corresponding Examples 4, 5, and 6A,B. A generalized scheme illustrating the mixture of products formed by denaturing and reannealing two such homoduplexes is provided in FIG. 5.

Detailed Description Paragraph Right (52):

Additional clonal types of all possible nucleotide substitutions at the polymorphic position, including 1, 2, 3, and 4 base pair insertion length variants were constructed using oligonucleotide cassette site-specific mutagenesis techniques. Briefly, short synthetic double stranded oligonucleotide inserts containing the desired nucleotide composition were subcloned into the unique Hpa I and Bgl II restriction sites which flank the polymorphic site. Each of the clonal allelic states was confirmed by conventional DNA sequencing. These clones provided precisely defined reagents for subsequent heteroduplex formation and detection. Desired polynucleotide fragments of variable length up to 1500 base pairs, each containing a single known polymorphic nucleotide, were generated by amplifying fragments from the appropriate plasmid clone using sets of PCR primers complementary to the vector sequence which flank the polymorphic locus at various defined distances.

Detailed Description Paragraph Right (54):

Double-stranded DNA homoduplex A, "homo-A-209", a 209-base pair fragment, was composed of two complementary 209-base fragments, polynucleotides 1 and 2. Double stranded DNA homoduplex G, "homo-G-209", a second 209-base pair fragment, was identical in sequence to homo-A-209 with the exception of one base pair (a G-C substituted for A-T present in homo-A-209) and was composed of polynucleotides 3 and 4. Polynucleotide 3 was identical in sequence to polynucleotide 1, with the exception of a guanosine at position 168 from the 5' end of polynucleotide 3, in comparison to an adenosine at the analogous position in polynucleotide 1. In a similar fashion, polynucleotide 4 was identical in sequence to polynucleotide 2, with the exception of a cytosine at position 42 from the 5' end replacing a thymidine in the same position in polynucleotide 2.

Detailed Description Paragraph Right (55):

The resulting mixture of products, containing original homoduplexes homo-A-209 and homo-G-209 and newly formed heteroduplexes hetero-AC-209 and hetero-GT-209 were then analyzed under the denaturing conditions of the present invention using ion pairing reverse phase HPLC. The designation hetero-AC-209 represents the double stranded product formed by annealing polynucleotides 1 and 4, and contains a single base pair A-C mismatch at position 168 relative to oligo 1. The designation hetero-GT-209 represents the double stranded product formed by annealing polynucleotides 2 and 3, and contains a single base pair G-T mismatch at position 168 relative to polynucleotide 3.

Detailed Description Paragraph Right (57):

At a less stringent column temperature of 50.degree. C. (FIG. 6C), separation/detection of the individual components of the product mixture containing homo-A-209, homo-G-209, hetero-AC-209, and hetero-GT-209 was not achieved (FIG. 6C). As seen in FIG. 6C, the components of the mixture co-eluted as a single peak, with a retention time about equivalent to that of each of homoduplexes homo-A-209 and homo-G-209 (FIGS. 6A,B).

Detailed Description Paragraph Right (58):

Upon raising the column temperature to 54.degree. C., the resolution of the separation was significantly enhanced as is shown in FIG. 7C. The two same-size homoduplex products were clearly separated both from each other and from the heteroduplexes, which eluted from the column slightly earlier than did the homoduplexes.

Detailed Description Paragraph Right (59):

Further to this point, FIGS. 10A-10C illustrate the sensitivity and resolving power of the method as a function of column temperature for samples of homo-A-209 and homo-G-209, subjected to denaturation and reannealing conditions, to produce a mixture containing homo-A-209, homo-G-209, hetero-AC-209, and hetero-GT-209. As seen in FIGS. 10A-C, at 50.degree. C. the mixture elutes as a single peak, with resolution improving at a heightened column temperature of 52.degree. C., and resulting in base line separation of the heteroduplexes from the homoduplex products at an optimized column temperature of 54.degree. C.

Detailed Description Paragraph Right (60):

In contrast to shorter double stranded DNA fragments having less than about 70 base pairs and containing a single base pair mismatch (e.g., Example 3), based on the number of peaks observed in FIG. 7C, larger DNA fragments appear to be only partially denatured using the optimized denaturing chromatographic conditions of the present invention, resulting in the formation of a "bubble" at the site of the base-pair mismatch. The distortion of the DNA duplex caused by this partial denaturation or bubble appears to cause a shift towards shorter retention times and allows the separation of heteroduplexes containing a single base pair mismatch from homoduplexes of the same size, as illustrated in FIG. 7C.

Detailed Description Paragraph Right (61):

The results described above show the effective separation of larger nucleic acid duplexes (i.e., over about 70 base pairs) containing a single base pair mismatch from homoduplexes of the same size by partial denaturation of the heteroduplexes using the denaturing HPLC conditions described herein, leading to shorter retention times.

Detailed Description Paragraph Right (63):

Summarizing the results, at a column temperature of 50.degree. C., each of the mixtures of homo- and heteroduplex products co-eluted and were detected as a single peak. At the lower column temperature, separation of the components of each of the reaction mixtures was not achieved, indicating the sensitivity of the separation method to temperature under the conditions employed.

Detailed Description Paragraph Right (64):

From the chromatograms shown in FIGS. 8C and 9C (column temperature of 56.degree. C.), it can be seen that the present method can be used for detecting single base pair mismatches in heteroduplexes up to at least 1 kilobase in size under chromatographic conditions similar to those used in the separation of shorter nucleic acid fragments, suggesting the general applicability of the method. Using the methods described herein, a single base pair mismatch was detected in polynucleotides 1500 base pairs in length, as illustrated in FIGS. 12A-12C. The results further suggest the potential for detecting single base pair mismatches in duplexes up to 2 kilobase in size.

Detailed Description Paragraph Right (65):

The method of the present invention can be used to detect heteroduplex molecules obtained by reverse transcription prior to amplification by PCR. The method of the present invention can also be extended to the quantification of competitive reverse transcriptase PCR (RT-PCR) reactions, namely via the detection of heteroduplex formation. In competitive RT-PCR, heteroduplex formation is one potential consequence of the homology between competitor and native products. The identification of heteroduplex RT-PCR products can be an important factor in determining the quantitative accuracy of the competitive assay results.

Detailed Description Paragraph Right (66):

Generally, in competitive RT-PCR, an RNA homolog (competitor) which contains the same sequence which is recognized by the reaction primers for the RNA of interest (native) is modified either by changing the presence of a restriction site or by altering the length of the sequence intervening between the primer binding sites to provide competitor mutant RNA. A range of known quantities of competitor RNA is added to multiple reactions each containing uniform amounts of the RNA preparation to be quantified (native). DNA expression is estimated by observing the relative amount of native and competitor products resulting from RT-PCR (Ferre, et al.; Volkenandt, et al.). The production of heteroduplexes during RT-PCR amplification can be confirmed using the denaturing HPLC method of the present invention.

Detailed Description Paragraph Right (69):

In utilizing such an approach, the presence of a peak(s) in the resulting chromatogram that does not correspond to a product observed by gel electrophoresis may correspond to a heteroduplex.

Detailed Description Paragraph Right (70):

To confirm whether or not a peak detected by denaturing HPLC corresponds to a heteroduplex, native and competitor RNA are reverse transcribed and amplified in separate reactions. The amplification products are then mixed together, followed by IP-RP-HPLC analysis of this mixture. The mixture is then heated (e.g., at 97.degree. C. for 3 minutes) followed by cooling to (e.g., to 4.degree. C.), followed by a second IP-RP-HPLC analysis. The appearance of a new product peak, and comparison of the retention times of any new peaks to those of any unaccounted peaks observed in the chromatograms of the initial set of RT-PCR reactions, indicates heteroduplex formation.

Detailed Description Paragraph Right (71):

In addition to identifying single base pair mismatches, the HPLC method of the invention can also be used for detecting indels in nucleic acid fragments. In exemplary experiments similar to those described above, a 440-mer homoduplex was mixed with a 439-mer homoduplex, identical to the 440-mer, with the exception of a single base pair deletion. Following denaturation and reannealing, the resulting mixture, containing both the original homoduplex molecules and the newly formed heteroduplexes, was chromatographed by the denaturing HPLC method of the invention (pH 7.0, 55.5.degree. C., 0.1M TEAA/0.1M TEAA in 25% acetonitrile) and heteroduplex formation

was observed. The heteroduplex indel molecules were separated from the homoduplex starting reagents in the resulting chromatogram and exhibited shorter retention times than those of the corresponding 439-mer and 440-mer homoduplexes. Using the present method, similar results have been obtained for indels having two, three, and four base pair deletions.

Detailed Description Paragraph Right (76):

In one embodiment, the mobile phase components are introduced into a mixer inside the column oven and mixed prior to contact with the sample. Alternatively, the mobile phase components may be mixed at ambient temperature and contacted with the sample injector, also maintained at ambient temperature outside of the column oven. Both of the above variations have been shown to be suitable for detection of heteroduplexes as has been described.

Detailed Description Paragraph Right (79):

In investigating the effect of column length and pre-column equilibration on the detection of 1000-mer heteroduplexes, heteroduplex formation was detected using a 60 cm length of tubing between the injector and the column, with a total length of 50 cm maintained at a column temperature of 56.degree. C. Upon increasing the length of tubing to 100 cm (with a total length of 90 cm maintained at an oven temperature of 56.degree. C.), improved separation of the heteroduplex species was observed.

Detailed Description Paragraph Right (80):

As has been discussed in section IIB above, one parameter which impacts the denaturing HPLC method of the present invention is pH. Generally, the pH of the mobile phase is maintained between about 7 and 9. In attempting to observe a single base mismatch in polynucleotides 1.5 kb in length, a preferred pH for carrying out the separation was found to be 7.5, as is described in Example 6B and further illustrated in FIGS. 12A-12C, and particularly in FIG. 12 C, which indicates heteroduplex detection in polynucleotide molecules containing up to 1500 base pairs.

Detailed Description Paragraph Right (82):

In investigating the effect of sample sequence on the temperatures effective for heteroduplex separation, the denaturing HPLC method of the invention was used to detect polymorphic sites contained within the 31-A-G fragment of the .beta.-globin gene (Myers, et al., 1986), where the region containing the polymorphism is flanked by GC-rich segments.

Detailed Description Paragraph Right (85):

The results of the above experiment indicate that although in most cases a temperature of 56.degree. C. is effective for heteroduplex separation, in some instances, and particularly for polynucleotides containing polymorphisms flanked by GC-rich regions, higher temperatures may be necessary.

Detailed Description Paragraph Right (89):

Exemplary chromatograms of DYS199 samples are shown in FIGS. 11A and 11B, illustrating HPLC traces of (i) a sample containing only homoduplex molecules derived from samples of African and Italian origin (FIG. 11A), and (ii) samples of African and native South American origin, indicating the presence of heteroduplexes. Of the samples examined, only samples of native South American origin exhibited the detected polymorphism.

Detailed Description Paragraph Right (93):

The heteroduplex separation and detection method of the present invention based on heteroduplex formation (e.g., of PCR products) is faster, simpler, more sensitive and more informative than the currently available procedures (such as RNase A cleavage mismatch). The denaturing HPLC method of the invention detects heteroduplex molecules in a mixture containing both heteroduplexes and homoduplexes by utilizing conditions effective to at least partially denature the heteroduplexes. Under such denaturing conditions, heteroduplexes exhibit slightly different retention times (typically shorter) from their homoduplex counterparts, thus providing a sensitive and convenient assay for detecting heteroduplex formation.

Detailed Description Paragraph Right (94):

Using the conditions described above, base pair mismatches and indels can be observed in heteroduplexes using the method of the present invention. The preferred size range

for these heteroduplexes ranges from approximately 30 to 1000 base pairs in length, although larger-sized heteroduplexes can be used as well. Heteroduplexes formed from molecules with mismatched nucleotides were detected in duplexes having a degree of divergence was as low as about 0.1%, and even lower. In addition, more complex mixtures of restriction fragments (e.g., 100-1000 bp size range) resulting from the post-PCR digestion of longer amplification products can be surveyed for the presence of heteroduplexes.

Detailed Description Paragraph Right (101):

The present invention provides a method for comparative DNA sequencing in which potentially all possible nucleotide mismatches and insertion/deletions within select amplified DNA fragments obtained from multiple animal or human subjects can be detected. In the context of the present invention, comparative DNA sequencing is carried out by amplifying DNA samples, typically up to at least about 1.5 kb in length, obtained from multiple subjects. The amplified DNA fragments are then surveyed, either individually or in pools containing up to about 10 samples, for the presence or absence of heteroduplexes using the denaturing high performance liquid chromatography method of the present invention. In surveying the samples, the amplified DNA fragments are denatured and allowed to reanneal. The resulting mixture of DNA fragments is then applied to a stationary reverse-phase support. The sample mixture is eluted with a mobile phase containing an ion-pairing reagent and an organic solvent. Sample elution is carried out under conditions effective to at least partially denature any heteroduplexes present in the sample and results in the detection of any heteroduplex molecules contained in the sample. The detection of a heteroduplex indicates the presence of a base pair mismatch and/or an insertion/deletion in the sample fragment(s).

Detailed Description Paragraph Right (102):

In instances in which only homoduplexes are observed during the sample screening, further standard sequencing is not required since the sequence is monomorphic (e.g., lacking polymorphic sites) in all subjects compared. In utilizing the method of the present invention, only those DNA fragments identified as heteroduplexes, and therefore identified as containing at least one polymorphic site, are then sequenced by conventional methods to characterize the observed polymorphism(s).

Detailed Description Paragraph Right (106):

Typically, probes for any target nucleic acid can be selected from a region of the microorganism's genomic material, such as rRNA (for example, as in Weisburg, et al.). In this way probes can be identified that will form homoduplexes to identify specific species. Formation of heteroduplexes indicates that the sequences have diverged from the probe sequence.

Detailed Description Paragraph Right (115):

Another embodiment of the present invention is the use of specific probes to identify variants based on the formation of homoduplex complexes. For example, sequences corresponding to a particular virus variant can be cloned and amplified. These cloned sequences are then used as a probe against viral molecules isolated from a number of test sources. Using the method of the present invention, if homoduplexes are formed in hybridization reactions between the probe and the test source, then the test source is shown to be similar to the cloned probe variant. If on the other hand heteroduplexes are formed between the probe and test sequences, then sequence divergence between the probe and test sequences is indicated.

Detailed Description Paragraph Right (116):

With respect to cancer, once a diagnosis has been made, and a region of DNA associated with the cancerous growth has been identified, the heteroduplex separation method of the present invention can be used to evaluate the extent of infiltration of tumor cells within a tissue population. Exemplary potential target sequences are protooncogenes, for example, including but not limited to the following: c-myc, c-myb, c-fos, c-kit, ras, and BCR/ABL (e.g., Gazdar, et al.; Wickstrom; Zalewski, et al.; Calabretta, et al., 1992, 1993;), oncogenes/tumor suppressor genes (e.g., p53, Bayever, et al.). In tumor cells, deletions, insertions, rearrangements and divergent sequences in such genes or in the regions of DNA surrounding the coding sequences of such genes, all allow formation of heteroduplexes between amplified variant DNA and amplified DNA from normal cells.

Detailed Description Paragraph Right (118):

Typically, samples to be analyzed by the method of the present invention are obtained by polymerase chain reaction amplification--the amplified sequences are denatured and reannealed before HPLC analysis. In addition to obtaining nucleic acid samples by amplification, other samples sources can be used as well. For example, sequences of interest can be cloned (e.g., in a lambda vector; Sambrook, et al.) from two different sources. The sequences of interest are independently isolated away from vector sequences (e.g., by restriction endonuclease digestion and fragment purification). These two samples can then be combined, denatured, renatured, and the resulting heteroduplexes analyzed in accordance with the present method.

Detailed Description Paragraph Right (145):

These results demonstrate the effect of increasing column temperature on the denaturation of short oligonucleotide homoduplexes with single stranded overhangs and further illustrate that such species (e.g., homoduplexes containing base overhangs) are detectable even under partially denaturing conditions (e.g., 40.degree. C.).

Detailed Description Paragraph Right (155):

Table 1 shows all possible nucleotide substitution heteroduplex mismatches formed from appropriate plasmid clones and subsequently detected by denaturing high performance liquid chromatography for exemplary size fragments up to 1.5 kb in length.

Detailed Description Paragraph Right (157):

Double-stranded DNA homoduplex A, "homo-A-209", a 209-base pair fragment was composed of two complementary 209-base fragments, polynucleotide 1 (SEQ ID NO:5) and polynucleotide 2 (SEQ ID NO:6). Double stranded DNA homoduplex G, "homo-G-209", a 209-base pair fragment identical in sequence to homo-A-209 with the exception of one base pair (a G-C substituted for A-T present in homo-A-209) was composed of polynucleotide 3 (SEQ ID NO:7) and complementary polynucleotide 4 (SEQ ID NO:8). Polynucleotide 3 was identical in sequence to polynucleotide 1, with the exception of a guanosine at position 168 from the 5' end of polynucleotide 3, in comparison to an adenosine at the analogous position in polynucleotide 1. In a similar fashion, polynucleotide 4 was identical in sequence to polynucleotide 2, with the exception of a cytosine at position 42 from the 5' end replacing a thymidine in the same position in polynucleotide 2.

Detailed Description Paragraph Right (158):

The double stranded oligonucleotides homo-A-209 and homo-G-209 were subjected to denaturation and reannealing under the following conditions in a Perkin Elmer 9600 thermal cycler: 95.degree. C. for 3 minutes, followed by cooling from 95.degree. C. to 65.degree. C. at a rate of 1.degree. C. per minute (e.g., over a period of 30 minutes), followed by sample storage at 6.degree. C. A schematic representation of the products formed by denaturing the above 209-mer homoduplexes followed by reannealing is provided in FIG. 5. The resulting mixture of products, containing original homoduplexes homo-A-209 and homo-G-209 and newly formed heteroduplexes hetero-AC-209 and hetero-GT-209 were then analyzed by IP-RP-HPLC. Hetero-AC-209 represents the double stranded product formed by annealing oligonucleotides 1 and 4, and contains a single base pair A-C mismatch at position 168 relative to oligo 1. Hetero-GT-209 represents the double stranded product formed by annealing oligonucleotides 2 and 3, and contains a single base pair G-T mismatch at position 168 relative to oligo 2.

Detailed Description Paragraph Right (159):

Samples of each of homo-A-209, homo-G-209, and the resulting heteroduplexes formed by denaturation and naturation of homo-A-209 and homo-G-209 were then directly chromatographed on alkylated nonporous PS-DVB particles (as described above) packed into a stainless steel column (NPC18M HPLC Biopolymer column, 50.times.4.6 mm I.D., Serasep, Inc., Santa Clara, Calif.). Eluent A consisted of 0.1M TEAA (triethylammonium acetate), pH 7.0 and Eluent B consisted of 0.1M TEAA, pH 7.0, in 25 % acetonitrile. The samples were eluted using a linear gradient profile of 37-63% Eluent B in 5.5 minutes, followed by 90% Eluent B for 1.5 minutes, at a flow rate of 1 ml per minute (UV detection, 254 nm). Two separate runs were performed at 50.degree. C. and 54.degree. C. to optimize the effect of column temperature on separation of the product mixture components, as shown in FIGS. 6A-6C and FIGS. 7A-7C, respectively.



Detailed Description Paragraph Right (160):

At a less stringent column temperature of 50.degree. C., separation/detection of the product mixture containing homo-A-209, homo-G-209, hetero-AC-209, and hetero-GT-209 was not achieved (FIG. 6C). However, upon raising the column temperature to 54.degree. C., the two homoduplex products were clearly separated from the heteroduplexes, which eluted from the column slightly faster than did the homoduplexes.

Detailed Description Paragraph Right (161):

In contrast to shorter double stranded DNA fragments having less than about 70 base pairs and containing a single base pair mismatch (e.g., Example 3), larger DNA fragments are only partially denatured using the optimized chromatographic conditions of the present invention, resulting in the formation of a "bubble" at the site of the base-pair mismatch. This partial denaturation or bubble causes a shift towards shorter retention times and allows the separation of heteroduplexes containing a single base pair mismatch from homoduplexes of the same size, as illustrated in FIG. 7C.

Detailed Description Paragraph Right (162):

Further to this point, FIGS. 10A-10C illustrate the sensitivity and resolving power of the method as a function of column temperature for samples of homo-A-209 and homo-G-209, subjected to denaturation and reannealing conditions, to produce a mixture containing homo-A-209, homo-G-209, hetero-AC-209, and hetero-GT-209. As seen in FIGS. 10A-C, at 50.degree. C. the mixture elutes as a single peak, with resolution improving at a heightened column temperature of 52.degree. C., and resulting in base line separation of the heteroduplexes from the homoduplex products at an optimized column temperature of 56.degree. C.

Detailed Description Paragraph Right (163):

In summary, the above results show the effective separation of larger nucleic acid duplexes (e.g. over 200 base pairs) containing a single base pair mismatch from homoduplexes of about the same size by partial denaturation of the heteroduplexes using the denaturing HPLC conditions described herein, leading to shorter retention times.

Detailed Description Paragraph Right (164):

In order to examine the sensitivity of the HPLC detection method of the invention, an experiment identical to that described in Example 4 above was carried out with two different DNA homoduplexes each containing 439 base pairs.

Detailed Description Paragraph Right (166):

The 439-base pair homoduplex A, designated as homo-A-439, was obtained by annealing polynucleotide 5 with polynucleotide 6. A 439-base pair homoduplex G, designated as homo-G-439, was obtained by annealing polynucleotide 7 with polynucleotide 8. Denaturation and reannealing were carried out as described above, generating a mixture containing homo-A-439, homo-G-439, and two resulting heteroduplex species.

Detailed Description Paragraph Right (167):

The resulting IP-RP-HPLC chromatograms for both homo-A-439 and homo-G-439 as well as for the sample obtained from denaturing and subsequent reannealing of homo-A-439 and homo-G-439 are shown in FIGS. 8A-8C, respectively. The column, column packing, composition of Eluent A and Eluent B, linear gradient, as well as the flow rate and detector wavelength are the same as those employed in Example 4. The separation was carried out at a column temperature of 56.degree. C. As seen in FIG. 8C, detection of the 439-mer heteroduplexes present in a sample also containing two 439-mer homoduplexes is achieved under the conditions employed, as evidenced by the peaks having shorter retention times and indicated by an arrow.

Detailed Description Paragraph Right (168):

The samples were also chromatographed at a column temperature of 50.degree. C., however, separation of the components of the reaction mixture containing both heteroduplex and homoduplex products was not achieved at the lower column temperature (not shown), since the products co-elute as a single peak.

Detailed Description Paragraph Right (169):

These results further support the general applicability of the method in separating and detecting single base pair mismatches in heteroduplexes containing nearly 500 base

pairs and further demonstrate the effect of column temperature in effecting at least partial denaturation of heteroduplex molecules in a sample containing both heteroduplexes and homoduplexes, as suggested by peaks (corresponding to the heteroduplex) having shorter retention times.

Detailed Description Paragraph Right (170):

To further investigate the separation capabilities of the present method, an experiment similar to that described in Example 5 above was carried out with two different DNA homoduplexes each 1 kilobase in length.

Detailed Description Paragraph Right (172):

The 1000-base pair homoduplex A, designated as homo-A-1kb, was obtained by annealing polynucleotide 9 with polynucleotide 10. 1000-base pair homoduplex G, designated as homo-G-1kb, was obtained in a similar manner. Denaturation and reannealing were carried out as described above to produce a sample containing homoduplexes homo-A-1kb and homo-G-1kb as well as the two heteroduplexes, hetero-AC-1kb and hetero-GT-1kb.

Detailed Description Paragraph Right (173):

The resulting IP-RP-HPLC chromatograms for both homo-A-1kb and homo-G-1kb as well as for the sample obtained from denaturing and subsequent reannealing of homo-A-1kb and homo-G-1kb are shown in FIGS. 9A-9C, respectively. The column, column packing, and composition of Eluent A and Eluent B were the same as those employed in Examples 4 and 5. A linear gradient of 40%-70% B in 7 minutes, followed by an increase to 90% B in 1.5 minutes was used to elute the products. A flow rate of 1 ml/minutes and a detector wavelength of 256 nm was employed. The separation was carried out at a column temperature of 56.degree. C. and resulted in the detection of 1-kilobase heteroduplexes containing only a single base pair mismatch in a sample also containing 1-kb homoduplexes.

Detailed Description Paragraph Right (174):

As in the previous examples, the samples were also chromatographed at a column temperature of 50.degree. C., however, separation/detection of the 1-kilobase heteroduplexes in a sample also containing 1-kilobase homoduplex products was not achieved at the lower column temperature (not shown). At the lower column temperature, the mixture of homo- and heteroduplex products co-elute and are detected as a single peak.

Detailed Description Paragraph Right (175):

The above results demonstrate the utility of the method in detecting single base pair mismatches in heteroduplexes up to 1 kilobase in size under chromatographic conditions similar to those used in the separation of shorter nucleic acid fragments, suggesting the general applicability of the method. The results further suggest the potential for detecting single base pair mismatches in duplexes larger than 1 kilobase in size.

Detailed Description Paragraph Right (176):

In a similar fashion to Example 6A described above, 1.5 kilobase heteroduplex molecules containing a single base pair mismatch were separated using the denaturing HPLC method of the present invention.

Detailed Description Paragraph Right (177):

The 1.5 kb fragment was amplified essentially as described above using primers SEQ ID NO:24 (forward) and SEQ ID NO:25 (reverse). The 1.5 kb A form sequence of polynucleotide 11 was annealed with polynucleotide 12 to create a 1.5 kb homoduplex, homo-A-1.5 kb. A similar 1.5 kb G form sequence was also generated from the complementary polynucleotides 13 and 14, homo-G-1.5 kb. Following mixing, denaturing and reannealing the 1.5 kb homo A and G duplexes, the single base mismatch of the resulting heteroduplex molecules was at position 829 from the 5' end of polynucleotides 11 and 13.

Detailed Description Paragraph Right (178):

The separation was effected at a pH of 7.5. The elution was carried out using a Gynkotek low-pressure gradient HPLC system (Gynkotek, Germering, Germany) at a temperature of 57.degree. C. and a flow rate of 1 ml/minute. A binary gradient system was employed (with components as described above), using a linear gradient profile of 62-71% Eluent B in 5 minutes. The chromatograms are shown in FIGS. 12A-12C, with FIG.

12C illustrating separation of the 1.5 kb heteroduplex molecules from the corresponding homoduplexes. The results suggest the potential for detecting single base mismatches in duplexes up to 2 kilobase in size using the present method.

Detailed Description Paragraph Right (186):

Exemplary chromatograms are shown in FIGS. 11A and 11B, illustrating HPLC traces of (i) a sample containing only homoduplex molecules derived from samples of Italian origin (FIG. 11A), (ii) in comparison with samples of native South American origin, indicating the presence of heteroduplexes.

Detailed Description Paragraph Left (8):

B. Denaturation/Reannealing of 209-mer Homoduplexes Formation and Detection of 209-mer Heteroduplexes

Detailed Description Paragraph Left (9):

B. Denaturation/Reannealing of 1500-Base Pair Homoduplexes: Formation and Detection of 1500-Base Pair Heteroduplexes

Detailed Description Paragraph Center (5):

IV. HPLC-Promoted Partial Denaturation and Detection of Large Heteroduplexes

Detailed Description Paragraph Center (20):

Denaturation/Reannealing of 439-mer Homoduplexes Formation and Detection of 439-mer Heteroduplexes

Detailed Description Paragraph Center (22):

A. Denaturation/Reannealing of 1000-Base Pair Homoduplexes: Formation and Detection of 1000-Base Pair Heteroduplexes

Detailed Description Paragraph Table (1):

TABLE 1	DUPLEX SPECIES	Clone #	A/T (1)	G/C (2)
T/A (3)	C/G (4)		A/T (1)	a/t A/C + G/T A/A + T/T
A/G + C/T	G/C (2)	G/T + A/C	g/c G/A + T/C	G/G + C/C T/A (3)
T/G + C/A	C/G (4)	C/T + A/G	C/C + G/G	C/A + T/G c/g

Lower case = homoduplexes Upper case = heteroduplexes

Other Reference Publication (8):

Oefner, P.J., et al., "High Resolution Liquid Chromatography of Fluorescent Dye-Labeled Nucleic Acids," Anal. Biochem. 223:39-46 (1994).

Other Reference Publication (13):

Nagamine et al, "A PCR Artifact: Generation of heteroduplexes", Am. J. Hum. Genet. 45:337-339, 1989.

Other Reference Publication (15):

Keen et al, "Rapid detection of single base mismatches as heteroduplexes on hydrolink gels", Trends Genet. 7(1):5, Jan. 1991.

CLAIMS:

1. A chromatographic method for separating heteroduplex and homoduplex DNA molecules in a mixture, comprising:

applying the mixture to a stationary reverse phase support,

eluting the heteroduplex and homoduplex molecules of said mixture with a mobile phase containing an ion-pairing reagent and an organic solvent, where said eluting is carried out under conditions effective to at least partially denature said heteroduplexes and where said eluting results in the separation of said heteroduplexes from said homoduplexes.

19. The method of claim 1, where prior to said applying step the DNA molecules are amplified using the polymerase chain reaction and the amplified DNA molecules denatured and renatured to form a mixture of heteroduplex and homoduplex DNA

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217

[Generate Collection](#)[Print](#)**Search Results - Record(s) 1 through 1 of 1 returned.**

- ☒ 1. 6265557. 09 May 97; 24 Jul 01. ABO histo-blood group O alleles of the baboon. Diamond; David, et al. 536/23.1; 435/320.1 536/23.2 536/23.5. C07H021/02 C07H021/04 C12N015/00.

[Generate Collection](#)[Print](#)

Term	Documents
ALLELE-SPECIFIC.DWPI,EPAB,JPAB,USPT.	1399
ALLELE-SPECIFICS	0
PROBES1	0
PROBE.DWPI,EPAB,JPAB,USPT.	211714
PROBEA.DWPI,EPAB,JPAB,USPT.	3
PROBEB.DWPI,EPAB,JPAB,USPT.	1
PROBEC.DWPI,EPAB,JPAB,USPT.	3
PROBED.DWPI,EPAB,JPAB,USPT.	12019
PROBEE.DWPI,EPAB,JPAB,USPT.	4
PROBEF.DWPI,EPAB,JPAB,USPT.	1
PROBEG.DWPI,EPAB,JPAB,USPT.	1
(ALLELE-SPECIFIC NEAR5 PROBES1 NEAR5 LABEL\$2 NEAR5 FLUORESCENT).USPT,JPAB,EPAB,DWPI.	1

[There are more results than shown above. Click here to view the entire set.](#)

[Previous Page](#)[Next Page](#)

**End of Result Set**

Generate Collection

L2: Entry 1 of 1

File: USPT

Jul 24, 2001

DOCUMENT-IDENTIFIER: US 6265557 B1

TITLE: ABO histo-blood group O alleles of the baboon

Detailed Description Paragraph Right (64):

Allele-specific PNA probes are constructed and are labeled with fluorescent markers such that the A-specific probe is red, the B-specific probe is blue, and the probe specific to the major (A-like) O allele is yellow. These labeled probes are used to determine the genotype of a biological sample. Assignment of fluorescence color for probes is arbitrary and is provided herein for convenience. A variety of fluorescence labels is known to those of ordinary skill in the art.